
Shortage of Bile Acid Sequestrants – Colestyramine Powder for Oral Suspension 4g (Questran) and Colestipol (Colestid)

Updated: 8th November 2019

Description of product affected

Colestyramine

- Colestyramine is licensed for use in the following indications^{1,2}:
 - Primary prevention of coronary heart disease in men between 35 and 59 years of age and with primary hypercholesterolaemia who have not responded to diet and other appropriate measures.
 - Reduction of plasma cholesterol in hypercholesterolaemia, particularly in those patients who have been diagnosed as Fredrickson's Type II (high plasma cholesterol with normal or slightly elevated triglycerides).
 - Relief of pruritus associated with partial biliary obstruction and primary biliary cirrhosis.
 - Relief of diarrhoea associated with ileal resection, Crohn's disease, vagotomy and diabetic vagal neuropathy.
 - Management of radiation-induced diarrhoea.
- The dose used varies between 4g and 36g daily according to the indication.^{1,2}
- For relief of diarrhoea, it is common practice to use colestyramine off-label to treat "bile acid diarrhoea" where considered clinically appropriate. In Cambridgeshire and Peterborough colestyramine is recommended first line for the management of bile acid malabsorption (unlicensed indication).
- Colestyramine is also used to help reduce the volume of jejunal and ileostomy outputs as a consequence of bowel resection for any cause³ and in patients with myeloma treatment induced diarrhoea that is unresponsive to loperamide.⁴

Colesevelam

- Colesevelam 625mg tablets⁵ has a much more restricted range of licensed indications limited to treatment of hypercholesterolaemia and is recommended in Cambridgeshire and Peterborough as second line therapy for bile acid malabsorption after colestyramine and where conventional anti-diarrhoea medication have failed.⁶

Colestipol

- Colestipol^{7,8} is restricted to treatment of hypercholesterolaemia and is **NON-FORMULARY** in Cambridgeshire and Peterborough.

Background

Colestyramine

- Bristol-Myers Squibb (BMS) divested the Questran range (Questran and Questran Light) to Cheplapharm at the end of July 2019.
- Questran sachets are currently unavailable with no further supplies expected until mid-2020.
- **Questran Light sachets are available in limited supplies through the wholesaler Alliance Healthcare.**
- **Generic colestyramine light 4g sachet (recently launched by Mylan), currently only available from the wholesaler, Phoenix.**

Colesevelam

- Cholestagel 625mg tablets (colesevelam): **currently available.**

Colestipol (non-formulary)

- Colestid Orange (colestipol) 5g granules: **currently available with Alliance Healthcare.**
- Colestid (colestipol) granules 5g (plain): currently out of stock until Mid-November 2019.

Alternative agents and management options

- Given the current issues with all of these agents it would seem prudent to keep patients on their existing treatment for as long as supplies remain available, provided they are benefitting from treatment.
- If there is a need to switch patients between treatments the following factors should be taken into consideration:
 - Both Questran and Questran Light sachets contain 4g anhydrous colestyramine per sachet.
 - Questran Light contains 30mg of aspartame per sachet (whereas Questran contains about 3.8g sugar) and this may lead to tolerability issues in some patients with IBS.⁶
 - In switching patients between colestipol (5g sachet) and colestyramine (4g sachet) or vice versa it would seem appropriate to switch on a sachet for sachet basis and then titrate according to response if needed (with specialist input if appropriate).
 - **There is no clear guidance on switching from colestyramine sachets to colesevelam (Cholestagel) 625mg tablets or vice versa. Advice from gastroenterology specialists locally has confirmed that an appropriate initiation dosage when switching adult patients from colestyramine to colesevelam would be:**
 - *Colesevelam (Cholestagel) taken as 625mg orally ONE to TWO times a day.*⁹

- *Patients will need to be monitored for their response to this treatment.*
- If prescribers have any concerns regarding individual patients, we would ask that you seek advice from the specialist that initiated the bile acid sequestrant in the first instance.
- Before switching medication, patient and carers as appropriate should be counselled about any change to their medication.

Pruritus

- For patients with pruritus due to liver or biliary problems who do not achieve adequate benefit from treatment with a bile acid sequestrant, there are a number of alternative treatments available.^{9,10}
- Specialist advice may be needed to ensure that symptom control is maintained.

Urgent Action

For GP practices:

- Questran sachets
 - GP practices should urgently identify patients prescribed Questran sachets to ensure patients have sufficient stock of their medication to last during the shortage period.
 - Where patients do not have sufficient supplies or stock is unavailable, the practices should encourage patients to liaise with pharmacies directly to ascertain their stock levels and which alternative product is available as different pharmacies use a range of wholesalers.
 - Where stock is unavailable, if clinically appropriate for the patient colestyramine light (generic or Questran Light) are available in limited supplies via the normal ordering process.
 - Where stock is unavailable, if clinically appropriate for the patient colestipol (Colestid) sachets may be ordered via the normal ordering process.
 - Where stock is unavailable, if clinically appropriate for the patient, colesevelam 625mg tablets may be ordered via the normal ordering process.
 - Prescriptions for Colestid (Colestipol) and Cholestagel (Colesevelam) 625mg tablets should be made on an acute basis and patients switched back to their normal product when their supplies resume.
 - Counsel the patient and their carers as appropriate about any change to their medication BEFORE any change is made.

For pharmacies:

- Support by checking your current stock levels of these products and informing your local practices.
- Support GPs in counselling patients regarding any change to their medication.

References

1. Bristol-Myers Squibb Pharmaceuticals limited. Questran Powder for Oral Suspension 4g. SPC, date of revision of the text, November 2017: <https://www.medicines.org.uk/emc/product/5715/smpc>
2. Bristol-Myers Squibb Pharmaceuticals limited. Questran Light. SPC, date of revision of the text, November 2017: <https://www.medicines.org.uk/emc/product/911/smpc>

3. Personal communication – Chair British Society of Gastroenterology Research Committee
4. Personal communication – Chief Pharmacist St Marks Hospital London and Haematology Pharmacist, Guys & St Thomas’ NHS Foundation Trust.
5. British National Formulary Issue 76 (Sept 2018- March 2019)
6. NICE. SeHCAT (tauroselcholic [75 selenium] acid) for the investigation of diarrhoea due to bile acid malabsorption in people with diarrhoea-predominant irritable bowel syndrome (IBS-D) or Crohn's disease without ileal resection. Diagnostics guidance [DG7]. Published Nov 2012. Available: <https://www.nice.org.uk/guidance/dg7>
7. Pfizer Limited. COLESTID Orange 5g. SPC, date of revision of the text, 02/2015: <https://www.medicines.org.uk/emc/product/129/smpc>
8. SANOFI. Cholestagel 625 mg film-coated tablets. SPC, date of revision of the text. 05 April 2017: <https://www.medicines.org.uk/emc/product/6142>
9. Personal communication – gastroenterologist at Cambridge Universities Hospital Foundation Trust.
10. Management of pruritis associated with cholestasis. UpToDate (subscription only)
11. Personal communication – liver specialist pharmacist Kings Hospital London

Acknowledgements

- Professor Ramesh P Arasaradnam, Consultant Gastroenterologist, Chair British Society of Gastroenterology Research Committee, Deputy Lead CRN (Gastroenterology) West Midlands
- Michelle O’Connor, Chair of BAD UK, Bile Acid Diarrhoea Support (<https://www.bad-uk.org/>)
- Scott Mercer, Specialist pharmacist rheumatology, Guy’s and St Thomas’ NHS Foundation Trust Specialist Liver Pharmacist, Kings College Hospital
- Chief Pharmacist, St Marks Hospital London North West University NHS Foundation Trust
- Haematology Pharmacist, Guy’s and St Thomas’ NHS Foundation Trust
- Gastroenterology Pharmacist, Guy’s and St Thomas’ NHS Foundation Trust.

Original document prepared by:

David Erskine, London and South East Regional Medicines Information, Guy’s and St Thomas’ NHS Foundation Trust, 31 May 2018, updated 11 June 2019: email: medicinesinformation@gstt.nhs.uk

Document modified by and for all correspondence please contact:

Cambridgeshire and Peterborough Clinical Commission Group, Medicines Optimisation Team. Published 26 June 2019, reviewed and amended 11th July 2019, further updated 5 September 2019; further updated 2nd October 2019 and 8th November 2019. Email: CAPCCG.prescribingpartnership@nhs.net.

Disclaimer: This memo can be adapted for local use. The content does not reflect national guidance. Some of this memo is based on **clinical opinion** from practitioners. Users should bear this in mind in deciding whether to base their policy on this document. Individual trusts should ensure that procedures for unlicensed medicines are followed where a foreign import drug is required in the interim. Any decision to prescribe off-label must take into account the relevant GMC guidance and NHS Trust governance procedures for unlicensed medicines. Prescribers are advised to pay particular attention to the risks associated with using unlicensed medicines or using a licensed medicine off-label. As with all prescribing, the prescriber is medically and legally responsible for the prescriptions they sign and for their decisions and actions when they supply and administer medicines or authorise or instruct others to do so.